

Acris Antibodies, Inc.

6815 Flanders Drive, Suite 140 San Diego, CA 92121 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

AM08107RP-N

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Monoclonal Antibody to CD8 - PE

Alternate names: CD8 alpha chain, CD8A, MAL, T-cell surface glycoprotein CD8 alpha chain, T-lymphocyte

differentiation antigen T8/Leu-2

Catalog No.: AM08107RP-N

Quantity: 0.1 mg
Concentration: 0.1 mg/ml

Background: The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that

mediates efficient cell to cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T cell receptor on the T lymphocyte recognize antigen displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The

functional coreceptor is either a homodimer composed of two alpha chains, or a

heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. CD8 is expressed on most thymocytes and approximately 1/3 of peripheral blood T cells in humans. CD8 alpha/beta heterodimers are expressed only on TCR alpha/beta T cells, whereas CD8 alpha

homodimers can be expressed on alpha/beta and gamma/delta T cells and some NK cells in humans. The pattern of expression in other species has not been well documented.

Uniprot ID: Q6R2U9

NCBI: NP 001001907

GenelD: 396627

Host / Isotype: Mouse / IgG2a

Clone: 76-2-11

OG/20130514

Format: State: Liquid purified Ig fraction.

Buffer System: PBS containing 0.09% Sodium Azide as preservative and a stabilizing

agent.

Label: PE - R-Phycoerythrin

Applications: Flow Cytometry: $\langle \cdot \rangle = 0.2 \,\mu\text{g}/10e6 \,\text{cells.}$ (Ref.1-3)

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody reacts with Porcine CD8 alpha (Mr. 32-34 kDa). (Ref-1-3)

Species: Pig.

Other species not tested.

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.



AM08107RP-N: Monoclonal Antibody to CD8 - PE

General Readings:

- 1. Pescovitz, M.D, .K. Lunney, and D.H. Sachs. 1984. J. Immunol. 133:368.
- 2. Yamada, K., D.H. Sachs, and H. DerSimonian. 1995. J. Immunol. 155:5249.
- 3. Smith, C.V., T. Sablinski, J.S. Arn, et al. 1994. J. Immunother. with Emphasis on Tumor Immunology 16:105.

