

Acris Antibodies, Inc.

6815 Flanders Drive, Suite 140 San Diego, CA 92121 UNITED STATES Phone: +1.858.888.7900

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com AM08111FC-N

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Monoclonal Antibody to CD5 - FITC

Alternate names: LEU-1, LEU1, Ly-1, Lymphocyte antigen T1/Leu-1, Lyt-1, T-cell surface glycoprotein CD5

Catalog No.: AM08111FC-N

Quantity: 0.5 mg
Concentration: 0.5 mg/ml

Background: In humans, CD5 is a 55kDa T lymphocyte single chain transmembrane glycoprotein. It is

present on all mature T lymphocytes, on most thymocytes and on many T cell leukemias and lymphomas. It reacts with a subpopulation of activated B cells. CD5/Lyt1 antigen is a monomeric type I transmembrane glycoprotein expressed on thymocytes, T lymphocytes, and a subset of B lymphocytes, but not on natural killer (NK) cells. It has been identified as the major ligand of the B cell antigen CD72. The frequency of CD5+ B cells exhibits strain dependent variation, and the phenotypic, anatomical, functional, developmental, and pathological characteristics of the CD5+ B cells suggest that they may represent a distinct lineage, known as B1 cells. Binding of CD5 on the T cell surface can augment alloantigen or mitogen induced lymphocyte proliferation and induces increased cytosolic free calcium, IL2 secretion, and IL2R expression. It has been proposed that CD5 negatively regulates

signal transduction mediated by the T cell and B cell receptors.

Host / Isotype: Mouse / IgG1

Clone: f43

Format: State: Liquid purified Ig fraction.

Buffer System: PBS containing 0.09% Sodium Azide as preservative.

Label: FITC - Fluorescein Isothiocyanate Isomer 1

Applications: Flow Cytometry: $\langle \ \ | = 1 \, \mu g / 10e6 \, \text{cells.}$ (Ref.1-2)

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: Specific to Feline CD5. (Mr. 67kDa)

This antibody is useful as a pan T Cell Marker. (Ref.1)

Species: Feline (Cat). Other species not tested.

Storage: Store the antibody undiluted at 2-8°C for one month or in (aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General Readings: 1. Ackley, C.D., and M.D. Cooper. 1992. Am. J. Vet. Res. 53:466-471.

2. Dean, G.A., S.L. Quackenbush, C.D. Ackley, M.D. Cooper, and E.A. Hoover. 1991. Vet.

Immunol. Immunopathol 28:327-335.

Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com