

Acris Antibodies, Inc.

6815 Flanders Drive, Suite 140 San Diego, CA 92121 UNITED STATES Phone: +1.858.888.7900

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

AM08146FC-N

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Monoclonal Antibody to Human IgM (Mu chain specific) - FITC

Alternate names: Human Immunoglobulin M

Catalog No.: AM08146FC-N

Quantity: 0.5 mg
Concentration: 0.5 mg/ml

Background: IgM normally constitutes about 10% of serum immunoglobulins. IgM antibody is prominent

in early immune responses to most antigens and is largely confined to plasma due to it's large size. Monomeric IgM is expressed as a membrane bound antibody on the surface of B

cells and as a pentamer when secreted by plasma cells.

Due to it's high valency IgM is more efficient than other isotypes is binding antigens with repeating epitopes (virus particles and red blood cells) and is more efficient than IgG in activiating the complement pathway. The gene for the mu constant region contains four

domains separated by short intervening sequences.

IgM measurement yields information about the body's immediate resistance and response to infection as well as information related to specific diseases. Decreased levels are associated with immune deficiency states, hereditary deficiencies, and myeloma. Increased levels can be associated with Waldenstrom's macroglobulinemia, chronic

infection and hepatocellular disease.

Host / Isotype: Mouse / IgG3

Clone: UHB

Format: State: Liquid purified Ig fraction.

Buffer System: PBS containing 0.09% Sodium Azide as preservative.

Label: FITC - Fluorescein Isothiocyanate Isomer 1

Applications: Indirect Immunofluorescent staining of IgM+ human B lymphocytes: $\langle \ \ | = 1 \, \mu g/10e6 \, \text{cells}.$

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody reacts with the heavy chain of human IgM as demonstrated by ELISA.

It may also react with IgM from other species.

Species: Human.

Other species not tested.

Storage: Store the antibody undiluted at 2-8°C for one month or in (aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com