

## Monoclonal Antibody to Bromodeoxyuridine (BrdU) - FITC

<b>Catalog No.:</b>	AM10021FC-N
<b>Quantity:</b>	0.2 mg
<b>Concentration:</b>	0.2 mg/ml
<b>Background:</b>	Bromodeoxyuridine (5-bromo-2-deoxyuridine, BrdU) is a synthetic nucleoside which is an analogue of thymidine. BrdU is commonly used in the detection of proliferating cells in living tissues. BrdU can be incorporated into the newly synthesized DNA of replicating cells (during the S phase of the cell cycle), substituting for thymidine during DNA replication. Antibodies specific to BrdU can then be used to detect the incorporated chemical; thus indicating cells that were actively replicating their DNA. Binding of the antibody requires denaturation of the DNA by heat or acid.
<b>Host / Isotype:</b>	Mouse / IgG1
<b>Clone:</b>	BRDU206
<b>Immunogen:</b>	BrdU conjugated to KLH. <b>Remarks:</b> Epitope: Bromodeoxyuridine.
<b>Format:</b>	<b>State:</b> Liquid purified Ig fraction. <b>Buffer System:</b> PBS, pH 7.4 containing 1% BSA as stabilizer and 0.05% Sodium Azide as preservative. <b>Label:</b> FITC
<b>Applications:</b>	Suitable for Immunohistochemistry and Immunocytochemistry (Frozen or Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections and cell smears) For IHC dilute concentrated antibody at 1/50-1/100, use streptavidin~biotin system or polymer system, incubate 30 minutes at room temperature. For staining of FFPE tissue sections, incubate with 4 N HCl for 20 minutes at RT, followed by treatment with proteolytic digestion for 10 minutes at RT. Immunofluorescence: 10-20 µg/ml (1/10-1/20), incubate for 2 hours in the dark at RT or it can also be incubated overnight at 4°C. Flow Cytometry: 0.2-1.0 µg/0.1 ml (1/200-1/1,000). Recommended Positive Control: Liver of experimental animal injected with BrdU or cell line grown in presence of BrdU. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This antibody recognises Bromodeoxyuridine (BrdU). Cellular Localization: Nuclear. <b>Species:</b> All. Other species not tested.

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)



- Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing. Shelf life: One year from despatch.
- General Readings:**
1. Williamson K et.al. Cytometry, 15: 162-168, 1994.
  2. Meyer JS et.al. Cancer, 71: 3531-3540, 1993 Patel, S et.al. Gut, 34:893-896, 1993.

---

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)