

## Polyclonal Antibody to Biotin - PE

**Alternate names:** Vitamin B4

**Catalog No.:** AP09204RP-N

**Quantity:** 1 ml

**Concentration:** 0.5 mg/ml

**Background:** Epitope tags are short peptide sequences that are easily recognized by tag-specific antibodies. Due to their small size, epitope tags do not affect the tagged protein's biochemical properties. Most often sequences encoding the epitope tag are included with target DNA at the time of cloning to produce fusion proteins containing the epitope tag sequence. This allows anti-epitope tag antibodies to serve as universal detection reagents for any tag containing protein produced by recombinant means. This means that anti-epitope tag antibodies are a useful alternative to generating specific antibodies to identify, immunoprecipitate or immunoaffinity purify a recombinant protein. The anti-epitope tag antibody is usually functional in a variety of antibody-dependent experimental procedures. Expression vectors producing epitope tag fusion proteins are available for a variety of host expression systems including bacteria, yeast, insect and mammalian cells. Rockland Immunochemicals produces anti-epitope tag antibodies against many common epitope tags including Myc, GST, GFP, 6X His, MBP, FLAG and HA. Rockland Immunochemicals also produces antibodies to other tags including FITC, Rhodamine (TRITC), DNP and biotin.

**Host / Isotype:** Goat / IgG

**Immunogen:** Biotin conjugated to Keyhole Limpet Hemocyanin (b-KLH)

**Format:** **State:** Liquid

**Purification:** Immunoaffinity chromatography using Biotin coupled to sepharose beads

**Buffer System:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml BSA as stabilizer and 0.05% (w/v) Sodium Azide as preservative

**Label:** PE – R-Phycoerythrin

*Absorption / Emission:* 490 nm, 545 nm and 565 nm / 580 nm

**Applications:** Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.

**Specificity:** Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, anti-Phycoerythrin, Biotin conjugated IgG and Biotin conjugated Bovine Serum Albumin.

**Storage:** Store the antibody undiluted at 2-8°C.  
DO NOT FREEZE!  
Shelf life: one year from despatch.

**For research and in vitro use only. Not for diagnostic or therapeutic work.**  
Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)