

Monoclonal Antibody to CD5 - FITC

Alternate names:	CD5, LEU1, Lymphocyte antigen T1/Leu-1, T-cell surface glycoprotein CD5
Catalog No.:	CL006F
Quantity:	0.1 mg
Concentration:	0.1 mg/ml
Background:	<p>In humans, CD5 is a 55kDa T lymphocyte single chain transmembrane glycoprotein. It is present on all mature T lymphocytes, on most thymocytes and on many T cell leukemias and lymphomas. It reacts with a subpopulation of activated B cells. CD5/Lyt1 antigen is a monomeric type I transmembrane glycoprotein expressed on thymocytes, T lymphocytes, and a subset of B lymphocytes, but not on natural killer (NK) cells. It has been identified as the major ligand of the B cell antigen CD72. The frequency of CD5+ B cells exhibits strain dependent variation, and the phenotypic, anatomical, functional, developmental, and pathological characteristics of the CD5+ B cells suggest that they may represent a distinct lineage, known as B1 cells. Binding of CD5 on the T cell surface can augment alloantigen or mitogen induced lymphocyte proliferation and induces increased cytosolic free calcium, IL2 secretion, and IL2R expression. It has been proposed that CD5 negatively regulates signal transduction mediated by the T cell and B cell receptors.</p>
Uniprot ID:	P13379
NCBI:	NP_031676.3
GeneID:	12507
Host / Isotype:	Mouse / IgG2b
Clone:	CG16
Immunogen:	C3H.CE - Ly 1.2 : DS from C3H spleen. Fusion Partner: Myeloma SP2/0 - Ag 14 (M5).
Format:	State: Liquid purified IgG fraction Purification: Protein G Chromatography Buffer System: PBS containing 0.02% Sodium Azide as preservative and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml Label: FITC
Applications:	Flow Cytometry Analysis (see Protocols). Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This Monoclonal antibody reacts with T cells from Mouse strains expressing the Ly 1.2 phenotype, but does not react with lymphocytes from mouse strains expressing the Ly 1.1 phenotype.
Species Reactivity:	Tested: Mouse.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

Protocols: **FLOW CYTOMETRY ANALYSIS:**

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2x10⁷ cells/ml in media A. Add 50 µl of this suspension to each tube (each tube will then contain 1 x 10⁶ cells, representing 1 test).
4. To each tube, add 0.1 - 0.2 µg* of this Ab per 10⁶ cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50 µl ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15 µl of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).

Results - Tissue Distribution

Mouse Strain: BALB/c
Cell Concentration: 1x10⁶ cells per tests
Antibody Concentration Used: 0.2 µg/10⁶ cells
Isotypic Control: FITC Mouse IgG2b

Results - Strain Distribution

Cell Concentration: 1x10⁶ cells per tests
Antibody Concentration Used: 0.2 µg/10⁶ cells
Strains Tested: AKR, ATH, BALB/c, CBA/J, C3H/He
Positive: AKR, ATH, BALB/c
Negative: CBA/J, C3H/He

Pictures:


<u>Cell Source</u>	<u>Percentage of cells stained above control:</u>
Thymus	98.9%
Spleen	33.5%
Lymph Node	88.6%
Bone Marrow	3.3%

FLOW CYTOMETRY ANALYSIS

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com



Cell Source: Thymus

Percentage of cells stained above control: 98.9 %

FLOW CYTOMETRY ANALYSIS - Tissue Distribution

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com