

Monoclonal Antibody to CD62L / L-Selectin - PE

Alternate names:	CD62 antigen-like family member L, LAM-1, LECAM1, LNHR, LYAM1, Leu-8, Leukocyte adhesion molecule 1, Leukocyte surface antigen Leu-8, Leukocyte-endothelial cell adhesion molecule 1, Lymph node homing receptor, SELL, TQ1, gp90-MEL
Catalog No.:	CL032RX
Quantity:	0.3 mg
Concentration:	0.1 mg/ml
Background:	L-selectin is expressed on most T and B lymphocytes, neutrophils, monocytes, eosinophils. ¹ Pre-incubation of lymphocytes with this antibody completely and specifically blocks binding of lymphocytes to high endothelial venules (HEV) in vitro ^{2,3,6} and the migration of lymphocytes to lymph nodes in vivo. ^{2,3} Polymorphonuclear cells preincubated with this antibody do not migrate to the inflammatory foci.
Uniprot ID:	P18337
NCBI:	NP_035476.1
GeneID:	20343
Host / Isotype:	Rat / IgG2a
Clone:	MEL-14
Immunogen:	Mouse B cell Lymphoma, 38C-14
Format:	State: Liquid purified Ig fraction Purification: Protein G Chromatography Buffer System: PBS containing 0.02% Sodium Azide as preservative and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml Label: PE
Applications:	Flow Cytometry. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This monoclonal antibody reacts with a 90 kDa protein which is involved with the homing of lymphocytes to peripheral lymph nodes.
Species Reactivity:	Tested: Mouse.
Storage:	Store the antibody undiluted at 2-8°C. DO NOT FREEZE! This product is photosensitive and should be protected from light. Shelf life: one year from despatch.
General References:	1) Fink, P., W. Gallatin, R. Reichert, et al. 1985. Homing receptor-bearing thymocytes, an immunocomponent cortical subpopulation. Nature 313: 233-235 2) Gallatin, W.M., I.L. Weissman., E.C. Butcher 1983. A cell surface molecule involved in

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

organ specific homing of lymphocytes. Nature 304:30-34

3) Lewinsohn, D.M., R.F. Bargatze, E.C. Butcher 1987. Leukocyte endothelial cell recognition: evidence of a common molecular mechanism shared by neutrophils, lymphocytes and other leukocytes. J. Immunology 138:4313-4321

4) Reichert, R., M. Gallitin, E. Butcher, et al.. 1984. A homing receptor bearing cortical thymocyte subset: Implications for thymus cell migration and the nature of cortisone-resistant thymocytes. Cell 38: 89-99

5) Siegelman, M., I.C. Cheng, I.L. Weissman, et al. 1990. The mouse lymph node homing receptor is identical with the lymphocyte cell surface receptor Ly-22: Role of the EGF domain in endothelial binding. Cell 61: 611-622

6) Jalkanen, S., R.F. Bargatze, J. Toyos, et al. 1987. Lymphocyte Recognition of High Endothelium: Antibodies to Distinct Epitopes of an 85-95-kD Glycoprotein Antigen Differentially Inhibit Lymphocyte Binding to Lymph Node, Mucosal, or Synovial Endothelial Cell. J. of Cell Biol. 105: 983-990

7) Goler M.L et al. 1997. T Cell Genetic Background Determines Maintenance of IL-12 Signaling. J. of Immunol. 159: 1767-1774

Protocols:

FLOW CYTOMETRY ANALYSIS:

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 0.5-1.0 μ g* of this Ab per 10^6 cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C. (It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50 μ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

Results - Tissue Distribution:

Mouse Strain: BALB/c

Cell Concentration: 1×10^6 cells per test

Antibody Concentration Used: 1.0 μ g/ 10^6 cells

Isotypic Control: PE Rat IgG2a

Cell Source: Percentage of cells stained above control:

Thymus: 89.1%

Spleen: 34.8%

Lymph Node: 88.5%

Results - Strain Distribution:

Cell Concentration: 1×10^6 cells per test


For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Antibody Concentration Used: 1.0 µg/10e6 cells
Strains Tested: BALB/c, CBA/J, C3H/He, C57BL/6, AKR
Positive: BALB/c, CBA/J, C3H/He, C57BL/6, AKR
Negative: none

Pictures:



Cell Source: Lymph Node
Percentage of cells stained above control: 88.5%

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com