

Monoclonal Antibody to CD72 (CD72.1 alloantigen) - FITC

Alternate names:	B-Cell marker, B-cell differentiation antigen CD72, Ly-32, Ly32, Lyb-2, Lymphocyte antigen 32
Catalog No.:	CL033F
Quantity:	0.1 mg
Concentration:	0.1 mg/ml
Background:	CD72 antigen is a member of the type II integral membrane glycoproteins which includes other related cell surface molecules such as the asialoglycoprotein receptors, CD23 and the Kupffer cell receptor. The function of CD72 is unknown but the exposure of B cells to CD72 antibodies activates a variety of signaling pathways and can induce MHC class II expression and B cell proliferation. CD72 antigen is expressed on all cells of B cell lineage with the exception of plasma cells and weakly on human tissue macrophages.
Uniprot ID:	P21855
NCBI:	XP_003086403.1
GeneID:	100504743
Host / Isotype:	Mouse / IgG2a
Clone:	CT-72.1
Format:	State: Liquid purified Ig fraction Buffer System: PBS, 0.09% sodium azide (NaN ₃) and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. Label: FITC
Applications:	Flow cytometry (For details please see "protocols" / "specificity" below). Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody reacts with the CD72 alloantigen CD72.1, a B-cell surface protein that is encoded by the Cd72a allele. CD72.1 is expressed on cells of the B cell lineage, except plasma cells ³ . Mouse strains expressing CD72.1 include C57L/-, C58/-, DBA/1, DBA/2, and SWR/J. Tissue Distribution by Flow Cytometry Analysis: (Representative Dot Plot) Mouse Strain Tested: DBA/2 Cell Concentration : 1x10 ⁶ cells per test Antibody Concentration Used: 0.25 µg/10 ⁶ cells Isotypic Control: FITC Mouse IgG2a. Species: Mouse. Other species not tested.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Storage:

Store the antibody at 2 - 8 °C for up to one month. For long term storage, aliquot and freeze unused portion at -20 °C in volumes appropriate for single usage. Avoid repeated freezing and thawing. This product is photosensitive and should be protected from light.

Shelf life: one year from despatch.

General References:

1. Robinson, W.H., Tutt Landolfi, M.M., Schafer, H. and Parnes, J.R. 1993 Biochemical identity of the mouse Ly-19.2 and Ly-32.2 alloantigens with the B cell differentiation antigen Lyb-2/CD72. *J. Immunol.* 151:4764-4772.
2. Luo, W., de Velde, H.V., Hoegen, I.V., Parnes, J.R. and Thielemans, K. 1992 Ly-1 (CD5), a membrane glycoprotein of mouse T lymphocytes and a subset of B cells, is a natural ligand of the B cell surface protein Lyb-2 (CD72). *J. Immunol.* 148:1630-1634.
3. Ying, H., J.I. Healy, C.C. Goodnow, and J.R. Parnes. 1998. Regulation of mouse CD72 gene expression during B lymphocyte development. *J. Immunol.* 161: 4760-4767.

Protocols:**FLOW CYTOMETRY ANALYSIS:****Method:**

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population.
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add ~ 0.25 μ g of antibody per 1×10^6 cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C. (It is recommended that the tubes are protected from light, since most fluorochromes are light sensitive.)
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50 μ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:


- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Pictures:



Lymphocytes FL1-H: Mouse anti mouse CD72.1, FL2-H: B222 PE

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com