

## Monoclonal Antibody to CD120a / TNFR1 - PE

<b>Alternate names:</b>	TNF-R1, TNF-RI, TNFR-I, Tnfrsf1a, Tumor necrosis factor receptor 1, Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor type I, p55, p60
<b>Catalog No.:</b>	CL044RT
<b>Quantity:</b>	25 Tests
<b>Background:</b>	<p>Tumor Necrosis Factor (TNF) is a cytokine whose function is mediated through two distinct cell surface receptors (TNF Receptor I and TNF Receptor II) that are included in the TNF Receptor superfamily along with FAS antigen and CD40. TNF Receptors I and II are 55 and 75 kDa members, respectively, of a family of cell surface molecules including nerve growth factor receptor, Fas/Apo1, CD30, OX40, and 41BB, which are characterized by cysteine rich motifs in the extracellular domain. While TNF Receptor I and TNF Receptor II share 28% sequence homology in the extracellular domains, their intracellular domains lack sequence homology, suggesting that they differ in their internal signal transduction pathways. TNF Receptor I contains an approximately 80 amino acid death domain near its carboxy terminus capable of transmitting an apoptotic signal through its interaction with TRADD (TNF Receptor I associated death domain protein), and subsequent interactions with FADD. TNF Receptor I can also activate the transcription factor NFkB via TRAF2 (TNF Receptor associated factor 2). The cytoplasmic domain of TNF Receptor I can directly interact with Jak kinase, thereby activating the JAK/STAT signal transduction cascade.</p> <p>TNF Receptor I is expressed by virtually all nucleated mammalian cells, including hepatocytes, monocytes and neutrophils, cardiac muscle cells, endothelial cells, and CD34 + hematopoietic progenitors. Both TNF alpha and TNF beta bind to TNF Receptor I.</p>
<b>Uniprot ID:</b>	<a href="#">P25118</a>
<b>NCBI:</b>	<a href="#">10090</a>
<b>Host / Isotype:</b>	Rat / IgG2a
<b>Clone:</b>	HM104
<b>Format:</b>	<p><b>State:</b> Liquid purified Ig fraction. <b>Purification:</b> Ion Exchange Chromatography. <b>Buffer System:</b> PBS containing 0.02% Sodium Azide as preservative and 0.5% BSA as stabilizer. <b>Label:</b> PE – R. Phycoerythrin (RPE)</p>
<b>Applications:</b>	<p>Flow Cytometry: Use 10 µl of neat antibody to label 10e6 cells in 100 µl. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.</p>
<b>Specificity:</b>	<p>This antibody recognises an extracellular region of mouse CD120a (TNFR1, p55). <b>Species:</b> Mouse. Other species not tested.</p>

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)

**Storage:**

Store the antibody undiluted at 2-8°C.

**DO NOT FREEZE!**

This product is photosensitive and should be protected from light.

Shelf life: One year from despatch.

---

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)