

Monoclonal Antibody to MHC Class II (I-Ek) - FITC

Alternate names:	MHC class II antigen I-Ek
Catalog No.:	CL074F
Quantity:	0.1 mg
Concentration:	0.1 mg/ml
Host / Isotype:	Mouse / IgG2a
Clone:	14-4-4S
Immunogen:	C3H Donor: C3H.SW Fusion Partner: SP2/0-Ag 14
Format:	State: Liquid purified Purification: Protein G Chromatography Buffer System: PBS, 0.02% NaN ₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. Label: FITC
Applications:	Flow Cytometry. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This monoclonal antibody is specific for cells expressing the Ia antigen coded for by the Ea subregion. The reaction pattern of this antibody with a panel of standard and recombinant haplotypes demonstrates that this antibody reacts with antigen Ia.m7, which is expressed by all haplotypes except b,f,q and s. This antibody can be used to quantitate or eliminate cells bearing the Ia.m7 antigen and is well suited for identifying Ia cell populations of positive mouse strains. Species: Mouse. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. This product is photosensitive and should be protected from light. Shelf life: one year from despatch.
General References:	1. Ozato, K., Mayer, N., and Sachs, D.H. 1980. Hybridoma cell line secreting monoclonal antibodies to mouse H-2 and Ia Antigens. J. Immunol.: 214:533-540.
Protocols:	<u>FLOW CYTOMETRY ANALYSIS:</u> Method: 1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M cell separation medium.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 0.2 - 0.5 μ g* of this Ab per 10^6 cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C. (It is recommended that the tubes are protected from light, since most fluochromes are light sensitive.)
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50 μ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

Results - Tissue Distribution by Flow Cytometry Analysis:

- Mouse Strain: CBA/J
Cell Concentration: 1×10^6 cells per tests
Antibody Concentration Used: 0.2 μ g/ 10^6 cells
Isotypic Control: FITC Mouse IgG2a

Cell Source Percentage of cells stained above control:

- Spleen: 53.5% see **FIGURE 1**
 Lymph Node: 26.6%
 Bone Marrow: 38.9%

Strain Distribution by Flow Cytometry Analysis:

- Procedure: As above
Antibody Concentration: 0.2 μ g/ 10^6 cells
Strains Tested: see **FIGURE 2**; For a more detailed strain distribution - see reference 1.

Pictures:

<u>Strain</u>	<u>H-2 Loci Alleles</u>								<u>+/-</u>
	<u>K</u>	<u>A_{β}</u>	<u>A_{α}</u>	<u>E_{β}</u>	<u>E_{α}</u>	<u>C4</u>	<u>C4S</u>	<u>D</u>	
A.TH	s	s	s	s	s	s	s	d	-
B10.A(3R)	b	b	b	b/k	k	d	d	d	+
AKR	k	k	k	k	k	k	k	k	+
C3H/He	k	k	k	k	k	k	k	k	+
A.TL	s	k	k	k	k	k	k	d	+
C57BL/6	b	b	b	b	b	b	b	b	-
BALB/c	d	d	d	d	d	d	d	d	+

Figure 2

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
 Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Figure 1

