

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com R1201TR Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



Polyclonal Antibody to Avidin - Texas Red

Alternate names:	AVD
	R1201TR
Catalog No.:	
Quantity:	20 mg
Concentration:	10.0 mg/ml (by UV absorbance at 280 nm)
Background:	The biological function of avidin is not known. It is a homotetrameric glycoprotein found in chicken egg white, and forms a strong non covalent specific complex with biotin (one molecule of biotin per subunit of avidin).
Uniprot ID:	<u>P02701</u>
NCBI:	<u>NP_990651.1</u>
GenelD:	<u>396260</u>
Host:	Rabbit
Immunogen:	Avidin [Hen Egg White].
Format:	 State: Lyophilized purified Ig fraction. Purification: Immunoaffinity chromatography. Buffer System: 0.01 M Sodium Phosphate, 0.14 M Sodium Chloride, pH 7.4 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. Label: Texas Red – (TM) Sulfonyl Chloride (Molecular Weight 625 daltons) Absorption / Emission: 596 nm / 620 nm Molar Ratio: 3.6 moles Texas Red(TM) per mole of Rabbit IgG. Reconstitution: Restore with 2.0 ml of deionized water (or equivalent).
Applications:	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum and anti-Avidin. No reaction was observed against Streptavidin.



R1201TR: Polyclonal Antibody to Avidin - Texas Red

Storage:Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then
aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear
after standing at room temperature.
This antibody is stable for one month at 2-8°C as an undiluted liquid.
Dilute only prior to immediate use.
Avoid repeated freezing and thawing.
Shelf life: One year from despatch.

General References: J.A. Titus, P.P. Haugland, S.D. Sharrow, D.M. Segal J. Immunol. Methods 50; 193, 1982. (Conjugation)