

## Polyclonal Antibody to Monkey IgG, IgA, IgM -FITC-

Alternate names: Monkey IgA, Monkey IgG, Monkey IgM

Catalog No.: R1250F Quantity: 1 mg

**Concentration:** 1.0 mg/ml (by UV absorbance at 280 nm)

Host: Goat

Immunogen: Monkey IgG, IgA, IgM whole molecules.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/ml

BSA (lgG and Protease free), 0.01% (w/v) Sodium Azide as preservative. Label: FITC – Fluorescein isothiocyanate (Molecular Weight 390 daltons)

Absorption / Emission: 495 nm / 528 nm Molar Ratio: 3.0 moles FITC per mole of Goat IgG.

**Reconstitution:** Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring lot-to-lot consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

**Specificity:** This product was prepared from monospecific antiserum by immunoaffinity

chromatography using antigens coupled to agarose beads.

Assay by immunoelectrophoresis resulted in a single precipitin arc against

anti-Fluorescein, anti-Goat Serum.

This reagent is suitable for the detection of all monkey isotypes and chain combinations.

Storage: Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: The and Feltkamp, Immunology 18; 865, 1970. (Conjugation)