

Polyclonal Antibody to Donkey IgG [H&L] -Texas Red (TM)-

Alternate names: Donkey Immunoglobulin G

Catalog No.: R1251TR

Quantity: 2 mg

Concentration: 2.0 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Donkey IgG whole molecule.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v)

Sodium azide as preservative.

Label: Texas Red – (TM) Sulfonyl Chloride (Molecular Weight 625 daltons)

Absorption / Emission: 596 nm / 620 nm

Molar Ratio: 2.8 moles Texas Red(TM) per mole of Rabbit IgG. **Reconstitution**: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring lot-to-lot consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Donkey IgG coupled to agarose beads.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit

Serum, Donkey IgG and Donkey Serum.

Storage: Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: J.A. Titus, P.P. Haugland, S.D. Sharrow, D.M. Segal J. Immunol. Methods 50; 193, 1982.

(Conjugation)