

Polyclonal Antibody to Rabbit IgG [H&L] -TRITC-

Alternate names: Rabbit Immunoglobulin G

Catalog No.: R1255T Quantity: 2 mg

Concentration: 2.0 mg/ml (by UV absorbance at 280 nm)

Host: Sheep

Immunogen: Rabbit IgG, whole molecule

Format: State: Lyophilized purified Ig fraction

Purification: Immunoaffinity Chromatography using Rabbit IgG coupled to agarose. **Buffer System:** 0.02M Potassium Phosphate, 0.15M Sodium Chloride, pH 7.2 containing 0.01% (w/v) Sodium Azide as preservative and 10 mg/ml BSA (IgG and Protease free) as

stabilizer.

Label: TRITC – Tetramethylrhodamine isothiocyanante (Molecular Weight 444 daltons)

Absorption / Emission: 550 nm / 570 nm

Molar Ratio: 2.9 moles TRITC per mole of Sheep IgG

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy and Flow Cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring lot-to-lot consistency.

Recommended Dilutions: FLISA: 1/10,000-1/50,000. Flow Cytometry: 1/500-1/2,500. IF Microscopy: 1/1,000-1/5,000.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: Assay by Immunoelectrophoresis resulted in a single precipitin arc against anti-Sheep

Serum, Rabbit IgG and Rabbit Serum.

Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: 1. J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. (Conjugation)