

Polyclonal antibody to Chicken IgG [H&L] - HRP

Alternate names: Chicken IgY
Catalog No.: R1300HRP
Quantity: 1.5 mg

Concentration: 1.5 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Chicken IgG whole molecule

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity Chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) Gentamicin Sulfate as

preservative.

Label: HRP - Horseradish Peroxidase

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunoblotting (Western or Dot blot), ELISA, Immunoperoxidase Electron

Microscopy and Immunohistochemistry as well as other phosphatase-antibody based

enzymatic assays requiring lot-to-lot consistency.

Recommended Dilutions: Western blot: 1/2,500-1/10,000.

ELISA: 1/145,000.

Immunohistochemistry: 1/1,000-1/5,000.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Chicken IgG coupled to agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase,

anti-Rabbit Serum, Chicken IgG and Chicken Serum.

Storage: Store vial at 4°C prior to restoration.

Restore with deionized water or equivalent; centrifuge product if not completely clear after

standing at room temperature. This product is stable for one month at 2-8°C as an undiluted liquid. For extended storage aliquot contents and freeze at -20°C or below.

Avoid cycles of freezing and thawing. Dilute only prior to immediate use. Shelf life: one year from despatch.

General References: 1. Farr & Nakane, J. Immunol Methods 47; 129-144, 1981. (Conjugation)