

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com R1301AP Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



## Polyclonal Antibody to Chicken IgG F(c) - AP-

Alternate names:	Chicken IgY
Catalog No.:	R1301AP
Quantity:	1 mg
Concentration:	0.4 mg/ml (by UV absorbance at 280 nm)
Host:	Rabbit
Immunogen:	Chicken IgG F(c) fragment.
Format:	<ul> <li>State: Liquid (sterile filtered) purified Ig fraction.</li> <li>Purification: Immunoaffinity chromatography.</li> <li>Buffer System: 0.05M Tris Chloride, 0.15M Sodium Chloride, 1mM Magnesium Chloride, 0.1mM Zinc Chloride, 50% (v/v) Glycerol; pH 8.0, containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) and 0.01% (w/v) and 0.01% Sodium Azide as preservative.</li> <li>Label: AP – Alkaline Phosphatase (Calf Intestine) (Molecular Weight 140,000 daltons)</li> </ul>
Applications:	Suitable for Immunoblotting (Western or Dot blot), ELISA and Immunohistochemistry as well as other phosphatase-antibody based enzymatic assays requiring lot-to-lot consistency. <u>Recommended Dilutions</u> : This product has been assayed against 1.0 µg of Chicken IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:9,000 of the reconstitution concentration is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Chicken IgG, Chicken IgG F(c) and Chicken Serum. No reaction was observed against Chicken IgG F(ab')2.
Storage:	Store the antibody undiluted at 2-8°C. <b>DO NOT FREEZE!</b> Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic activity. Dilute only prior to immediate use. Shelf life: one year from despatch.
General References	: Modified from Avarameas and Ternyrock, Immunochemistry 32: 1175 1971. (Conjugation)

General References: Modified from Avarameas and Ternyrock, Immunochemistry 32; 1175 1971. (Conjugation)