

Polyclonal Antibody to Chicken IgG F(ab')2 - Texas Red (TM)-

Alternate names: Chicken IgY
Catalog No.: R1302TR
Quantity: 1.5 mg

Concentration: 1.5 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Chicken IgG F(ab')2 fragment.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v)

sodium azide as preservative.

Label: Texas Red – (TM) Sulfonyl Chloride (Molecular Weight 625 daltons)

Absorption / Emission: 596 nm / 620 nm

Molar Ratio: 3.1 moles Texas Red per mole of Rabbit IgG.

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring lot-to-lot consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Chicken IgG coupled to agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit

Serum, Chicken IgG, Chicken IgG F(ab') 2, and Chicken Serum.

No reaction was observed against Chicken IgG F(c).

Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: J. Titus, P. Haugland, S. Sharrow, D. Segal J. Immunol. Methods 50; 193, 1982.

(Conjugation)