

## Polyclonal Antibody to Chicken IgG [H&L] (multi-species ads) -Texas Red

|                            |  |
|----------------------------|--|
| <b>Alternate names:</b>    | Chicken IgY  |
| <b>Catalog No.:</b>        | R1303TR  |
| <b>Quantity:</b>           | 1 mg   |
| <b>Concentration:</b>      | 1.0 mg/ml (OD280)  |
| <b>Host:</b>               | Goat   |
| <b>Immunogen:</b>          | Chicken IgG whole molecule.  |
| <b>Format:</b>             | <b>State:</b> Lyophilized purified Ig fraction.<br><b>Purification:</b> Immunoaffinity chromatography.<br><b>Buffer System:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, with 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% Sodium Azide as a preservative.<br><b>Label:</b> Texas Red – (TM) Sulfonyl Chloride (Molecular Weight 625 daltons)<br><i>Absorption / Emission:</i> 596 nm / 620 nm<br><i>Molar Ratio:</i> 4.5 moles Texas Red per mole of Goat IgG.<br><b>Reconstitution:</b> Restore with 1.0 ml of deionized water (or equivalent). |
| <b>Applications:</b>       | Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.  |
| <b>Specificity:</b>        | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Chicken IgG and Chicken Serum. No reaction was observed against Bovine, Goat, Hamster, Horse, Human, Mouse, Rabbit, Rat and Sheep Serum Proteins.   |
| <b>Storage:</b>            | Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.  |
| <b>General References:</b> | 1. J.A. Titus, et al., J. Immunol. Methods 50; 193, 1982. (Conjugation)  |

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)