

Polyclonal Antibody to Bovine IgG (H+L chain) - FITC

Alternate names: Immunoglobulin G

Catalog No.: R1304F Quantity: 2 mg

Concentration: 2.0 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Bovine IgG, whole molecule

Format: State: Lyophilized purified IgG fraction

Purification: Immunoaffinity chromatography using Bovine IgG coupled to agarose beads

followed by solid phase adsorptions to remove any unwanted reactivities

Buffer System: 0.01 M Sodium Phosphate, 0.14 M Sodium Chloride, pH 7.4, with 0.01% (w/v) Thimerisol as preservative and 10 mg/ml Polyethylene Glycol (PEG mw 8,000) as

stabilizer

Label: FITC – Diaminotriazinylaminofluorescein (DTAF) (Molecular Weight 530 daltons)

which has identical fluorescence properties as Absorption / Emission: 495 nm / 528 nm

Molar Ratio: 4.2 moles DTAF per mole of Rabbit IgG

Reconstitution: Restore with 1.0 ml of deionized water or equivalent.

Applications: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring lot-to-lot consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: Assay by immunoelectrophoresis resulted in a single precipitin arc against

anti-Fluorescein, anti-Rabbit Serum, Bovine IgG, Bovine IgG and Bovine Serum.

Storage: Store vial at 2-8°C prior to restoration.

Restore with deionized water (or equivalent); centrifuge product if not completely clear after standing at room temperature. This product is stable for one month at 2-8°C as an

undiluted liquid.

For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Avoid repeated freezing and thawing.

Dilute only prior to immediate use. Shelf life: one year from despatch.

General References: Blakeslee & Baines, J. Immunol. Methods 13; 305-320, 1976. (Conjugation)