

Polyclonal Antibody to Bovine IgG (F(ab)2 specific) - FITC

Alternate names: Immunoglobulin G

Catalog No.: R1306F Quantity: 2 mg

Concentration: 2.0 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Bovine IgG F(ab')2 fragment.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.01 M Sodium Phosphate, 0.14 M Sodium Chloride, pH 7.4 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) sodium

azide as preservative.

Label: FITC – Diaminotriazinylaminofluorescein (DTAF) (Molecular Weight 530 daltons)

Absorption / Emission: 495 nm / 528 nm

Molar Ratio: 1.9 moles DTAF per mole of Rabbit IgG.

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring lot-to-lot consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Bovine IgG coupled to agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities.

Assay by immunoelectrophoresis resulted in a single precipitin arc against

anti-Fluorescein, anti-Rabbit Serum, Bovine IgG, Bovine IgG F(ab')2 and Bovine Serum.

No reaction was observed against Bovine IgG F(c).

Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: Blakeslee & Baines, J. Immunol. Methods 13; 305-320, 1976. (Conjugation)