

Polyclonal Antibody to Cat IgG F(c) -FITC-

Alternate names: Cat IgG Fc, Immunoglobulin G Fc

Catalog No.: R1308F
Quantity: 2 mg

Concentration: 2.0 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Cat IgG F(c) fragment.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v)

sodium azide as preservative.

Label: FITC – Fluorescein isothiocyanate (Molecular Weight 390 daltons)

Absorption / Emission: 495 nm / 528 nm

Molar Ratio: 3.5 moles FITC per mole of Rabbit IgG.

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring lot-to-lot consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Cat IgG coupled to agarose beads.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Cat IgG, Cat IgG F(c) and Cat Serum.

No reaction was observed against Cat IgG F(ab')2.

Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: The and Feltkamp, Immunology 18; 865, 1970. (Conjugation)