

Polyclonal Antibody to Cat IgG F(ab')2 -TRITC-

Alternate names: Cat IgG, Immunoglobulin G

Catalog No.: R1309T Quantity: 2 mg

Concentration: 2.0 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Cat IgG F(ab')2 fragment.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with Polyethylene Glycol (PEG-8000) as stabilizer and 0.01% (w/v) sodium azide as

preservative.

Label: TRITC – Tetramethylrhodamine isothiocyanante (Molecular Weight 444 daltons)

Absorption / Emission: 550 nm / 570 nm

Molar Ratio: 2.3 moles TRITC per mole of Rabbit IgG.

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy (1/500-1/2,500) and Flow cytometry (1/2,000-1/10,000) or

FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot

consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Cat IgG coupled to agarose beads.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti- Rabbit

Serum, Cat IgG, Cat IgG F(ab')2 and Cat Serum. No reaction was observed against Cat IgG F(c).

Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. (Conjugation)