

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com R1310F Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



## Polyclonal Antibody to Dog IgG [H&L] -DTAF-

Alternate names:	Dog IgG, Dog Immunoglobulin G
Catalog No.:	R1310F
Quantity:	2 mg
Concentration:	2.0 mg/ml (by UV absorbance at 280 nm)
Host:	Rabbit
Immunogen:	Dog IgG whole molecule
Format:	<ul> <li>State: Lyophilized purified Ig fraction</li> <li>Purification: Immunoaffinity chromatography</li> <li>Buffer System: 0.01 M Sodium Phosphate, 0.14 M Sodium Chloride, pH 7.4 with 10 mg/ml</li> <li>Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v)</li> <li>Thimerisol as preservative</li> <li>Label: FITC – Diaminotriazinylaminofluorescein (DTAF) (Molecular Weight 530 daltons)</li> <li>Absorption / Emission: 495 nm / 528 nm</li> <li>Molar Ratio: 2.4 moles DTAF per mole of Rabbit IgG</li> <li>Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).</li> </ul>
Applications:	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Dog IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Dog IgG and Dog Serum.
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: Blakeslee & Baines, J. Immunol. Methods 13; 305-320, 1976. (Conjugation)