

Polyclonal Antibody to Dog IgG [H&L]-Texas Red (TM)-

Alternate names: Dog IgG, Dog Immunoglobulin G

Catalog No.: R1310TR
Quantity: 2 mg

Concentration: 2.0 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Dog IgG whole molecule.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as

preservative.

Label: Texas Red – (TM) Sulfonyl Chloride (Molecular Weight 625 daltons)

Absorption / Emission: 596 nm / 620 nm

Molar Ratio: 3.6 moles Texas Red per mole of Rabbit IgG.

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy (1(/500-1/2,500) and Flow cytometry (1/2,000-1/10,000) or

FACS analysis as well as other antibody base fluorescent assays requiring lot-to-lot

consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Dog IgG coupled to agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit

Serum, Dog IgG, and Dog Serum.

Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: J. Titus, P. Haugland, S. Sharrow, D. Segal J. Immunol. Methods 50; 193, 1982.

(Conjugation)