

Polyclonal Antibody to Goat IgG (H+L chain) - TRITC

Alternate names: Goat Immunoglobulin G

Catalog No.: R1317T Quantity: 2 mg

Concentration: 2.0 mg/mL (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Goat IgG, whole molecule

Format: State: Lyophilized purified Ig fraction

Purification: Immunoaffinity Chromatography

Buffer System: 0.02M Potassium Phosphate, 0.15M Sodium Chloride, pH 7.2

Preservatives: 0.01% (w/v) Sodium Azide

Stabilizers: 10 mg/ml BSA (IgG and Protease free)

Label: TRITC – Tetramethylrhodamine isothiocyanante, Molecular Weight 444 daltons)

Absorption / Emission: 550 nm / 570 nm

Molar Ratio: 1.9 moles Rhodamine per mole of IgG.

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy (1/1,000-1/5,000) and Flow Cytometry (1/500-1/2,500) or

FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot

consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by Immunoaffinity

Chromatography using Goat IgG coupled to agarose beads.

Assay by Immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit

Serum, Goat IgG and Goat Serum.

Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. (Conjugation)