

Polyclonal Antibody to Goat IgG F(c) -AP-

Alternate names: Goat Immunoglobulin G

Catalog No.: R1319AP

Quantity: 1 mg

Concentration: 0.73 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Goat IgG F(c) fragment.

Format: State: Liquid (sterile filtered) purified Ig fraction

Purification: Immunoaffinity chromatography.

Buffer System: 0.05M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. **Label:** AP – Alkaline Phosphatase (Calf Intestine) (Molecular Weight 140,000 daltons)

Applications: Suitable for Immunoblotting (Western or Dot blot), ELISA and Immunohistochemistry as

well as other phosphatase-antibody based enzymatic assays requiring lot-to-lot

consistency.

Recommended dilutions: This product has been assayed against 1.0 ug of Goat IgG in a standard capture ELISA using pNPP (p-nitrophenyl phosphate) as a substrate for 30 minutes at room temperature. A working dilution of 1:400 to 1:2,800 of the reconstitution

concentration is suggested for this product.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Goat IgG coupled to agarose beads.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Goat IgG, Goat IgG F(c) and Goat Serum.

No reaction was observed against Goat IgG F(ab').

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic

activity.

Dilute only prior to immediate use. Shelf life: one year from despatch.

General References: Modified from Avarameas and Ternyrock, Immunochemistry 32; 1175 1971. (Conjugation)