

## Polyclonal Antibody to Goat IgG F(ab')<sub>2</sub> -AP-

<b>Alternate names:</b>	Goat Immunoglobulin G
<b>Catalog No.:</b>	R1320AP
<b>Quantity:</b>	1 mg
<b>Concentration:</b>	1.0 mg/ml (by UV absorbance at 280 nm)
<b>Host:</b>	Rabbit
<b>Immunogen:</b>	Goat IgG F(ab') <sub>2</sub> fragment.
<b>Format:</b>	<b>State:</b> Liquid (sterile filtered) purified Ig fraction. <b>Purification:</b> Immunoaffinity chromatography. <b>Buffer System:</b> 0.05M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. <b>Label:</b> AP – Alkaline Phosphatase (Calf Intestine) (Molecular Weight 140,000 daltons)
<b>Applications:</b>	Suitable for Immunoblotting (Western or Dot blot), ELISA and Immunohistochemistry as well as other phosphatase-antibody based enzymatic assays requiring lot-to-lot consistency. <b>Recommended Dilutions:</b> This product has been assayed against 1.0 ug of Goat IgG in a standard capture ELISA using pNPP (p-nitrophenyl phosphate) as a substrate for 30 minutes at room temperature. A working dilution of 1:500 to 1:3,000 is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Goat IgG, Goat IgG F(ab') <sub>2</sub> and Goat Serum. No reaction was observed against Goat IgG F(c).
<b>Storage:</b>	Store the antibody undiluted at 2-8°C. <b>DO NOT FREEZE!</b> Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic activity. Dilute only prior to immediate use. Shelf life: one year from despatch.
<b>General References:</b>	(Conjugation): Modified from Avrameas and Ternyrock, <i>Immunochemistry</i> 32; 1175 1971.

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)