

## Polyclonal Antibody to Goat IgG F(ab')<sub>2</sub> -HRP-

<b>Alternate names:</b>	Goat Immunoglobulin G
<b>Catalog No.:</b>	R1320HRP
<b>Quantity:</b>	2 mg
<b>Concentration:</b>	2.0 mg/ml (by UV absorbance at 280 nm)
<b>Host:</b>	Rabbit
<b>Immunogen:</b>	Goat IgG F(ab') <sub>2</sub> fragment.
<b>Format:</b>	<b>State:</b> Lyophilized purified IgG fraction. <b>Purification:</b> Immunoaffinity chromatography. <b>Buffer System:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Gentamicin Sulfate as preservative. <b>Label:</b> HRP – Horseradish Peroxidase <b>Reconstitution:</b> Restore with 1.0 ml of deionized water (or equivalent).
<b>Applications:</b>	Suitable for Immunoblotting (Western or Dot blot), ELISA, Immunoperoxidase electron microscopy and Immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency. <b>Recommended Dilutions:</b> This product has been assayed against 1.0 µg of Goat IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:7,000 to 1:35,000 of the reconstitution concentration is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Goat IgG, Goat IgG F(ab') <sub>2</sub> and Goat Serum. No reaction was observed against Goat IgG F(c).
<b>Storage:</b>	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.
<b>Caution:</b>	Do <u>Not</u> Use Sodium Azide as Preservative.
<b>General References:</b>	Farr & Nakane, J. Immunol. Methods 47; 129-144. 1981 (Conjugation)

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)