

Polyclonal Antibody to Hamster IgG [H&L] -FITC-

Catalog No.: R1327F

Quantity: 1.5 mg

Concentration: 1.0 mg/ml

Host: Rabbit

Immunogen: Hamster IgG whole molecule.

Format: State: Lyophilized purified Ig fraction

Purification: Immunoaffinity Chromatography using Hamster IgG coupled to agarose. **Buffer System:** 0.01 M Sodium Phosphate, 0.14 M Sodium Chloride, pH 7.4 containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v)

Sodium azide as preservative.

Label: FITC - Flourescence isothiocyanate, Mol Weight 390 daltons

Absorption / Emission: 495 nm / 528 nm

Molar Ratio: 4.5 moles FITC per mole of Rabbit IgG

Reconstitution: Restore with 1.5 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring lot-to-lot consistency.

Recommended Dilution(s):

Flow Cytometry: 1/2,000-1/10,000. IF Microscopy: 1/500-1/2,500.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: Assay by immunoelectrophoresis resulted in a single precipitin arc against

anti-Fluorescein, anti-Rabbit Serum, Hamster IgG and Hamster Serum.

Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: 1. Blakeslee & Baines, J. Immunol. Methods 13; 305-320, 1976. (Conjugation)