

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com R1331TR Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



## Polyclonal Antibody to Horse IgG F(ab')2 -Texas Red (TM)

Alternate names:	Horse IgG, Horse Immunoglobulin G
Catalog No.:	R1331TR
Quantity:	1.5 mg
Concentration:	1.5 mg/ml (by UV absorbance at 280 nm)
Host:	Rabbit
Immunogen:	Horse IgG F(ab')2 fragment.
Format:	<ul> <li>State: Lyophilized purified Ig fraction.</li> <li>Purification: Immunoaffinity chromatography.</li> <li>Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative.</li> <li>Label: Texas Red – (TM) Sulfonyl Chloride (TR; Molecular Weight 625 daltons) Absorption / Emission: 596 nm / 620 nm Molar Ratio: 3.0 moles Texas Red per mole of Rabbit IgG.</li> <li>Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).</li> </ul>
Applications:	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Horse IgG, Horse IgG F(ab')2 and Horse Serum. No reaction was observed against Horse IgG F(c).
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.
Conoral Deferences	A Titue at al Limmunal Matheda 50, 102, 1082 (Canivastian)

General References: J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. (Conjugation)

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.