

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

R1332T Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



Polyclonal Antibody to Human IgG [H&L] -TRITC-

| Alternate names: | Human Immunoglobulin G |
|--------------------|--|
| Catalog No.: | R1332T |
| Quantity: | 1 mg |
| Concentration: | 1.0 mg/ml (by UV absorbance at 280 nm) |
| Host: | Chicken |
| Immunogen: | Human IgG whole molecule |
| Format: | State: Lyophilized purified Ig fraction. Purification: Immunoaffinity Chromatography. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: TRITC – Tetramethylrhodamine isothiocyanante ; Molecular Weight 444 daltons) Absorption / Emission: 550 nm / 570 nm Molar Ratio: 3.3 moles TRITC per mole of Chicken IgG Reconstitution: Restore with 1.0 ml of deionized water (or equivalent). |
| Applications: | Suitable for Immunomicroscopy and Flow Cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. <u>Recommended Dilutions:</u> FLISA: 1/10,000-1/50,000. IF Microscopy: 1/1,000-1/5,000. Flow Cytometry: 17500-1/2,500. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user. |
| Specificity: | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Chicken Serum, Human IgG and Human Serum. |
| Storage: | Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch. |
| General References | : I.A. Titus, et al. I. Immunol, Methods 50: 193, 1982. (Conjugation) |

General References: J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. (Conjugation)

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.