

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

## R1334F Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



## Polyclonal Antibody to Human IgG [H&L] -DTAF-

| Alternate names:   | Human Immunoglobulin G   |
|--------------------|--|
| Catalog No.:       | R1334F   |
| Quantity:          | 2 mg   |
| Concentration:     | 2.0 mg/ml (by UV absorbance at 280 nm)   |
| Host:              | Rabbit   |
| Immunogen:         | Human IgG, whole molecule.   |
| Format:            | <ul> <li>State: Lyophilized purified Ig fraction.</li> <li>Purification: Immunoaffinity chromatography.</li> <li>Buffer System: 0.01 M Potassium Phosphate, 0.14 M Sodium Chloride, pH 7.6 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Thimerosal as preservative.</li> <li>Label: FITC – Dichlorotriazinylaminofluorescein Dihydrochloride (DTAF) (Molecular Weight 530 Daltons)</li> <li>Absorption / Emission: 495 nm / 528 nm</li> <li>Molar Ratio: 3.6 moles DTAF per mole of Rabbit IgG.</li> <li>Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).</li> </ul> |
| Applications:      | Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other<br>antibody based fluorescent assays requiring lot-to-lot consistency.<br>Other applications not tested. Optimal dilutions are dependent on conditions and should<br>be determined by the user.   |
| Specificity:       | This product was prepared from monospecific antiserum by immunoaffinity<br>chromatography using Human IgG coupled to agarose beads followed by solid phase<br>adsorption(s) to remove any unwanted reactivities.<br>Assay by immunoelectrophoresis resulted in a single precipitin arc against<br>anti-Fluorescein, anti-Rabbit Serum, Human IgG and Human Serum.<br>No reaction was observed against Mouse Serum Proteins.  |
| Storage:           | Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then<br>aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear<br>after standing at room temperature.<br>This antibody is stable for one month at 2-8°C as an undiluted liquid.<br>Dilute only prior to immediate use.<br>Avoid repeated freezing and thawing.<br>Shelf life: One year from despatch.  |
| General References | : The and Feltkamp, Immunology 18; 865, 1970. (Conjugation)  |

General References: The and Feltkamp, Immunology 18; 865, 1970. (Conjugation)