

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com R1340F Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



## Polyclonal Antibody to Human IgM mu (mu chain specific) -FITC-

Alternate names:	Human Immunoglobulin M
Catalog No.:	R1340F
Quantity:	1 mg
Concentration:	1.0 mg/ml (by UV absorbance at 280 nm)
Host:	Goat
Immunogen:	Human IgM m mu chain.
Format:	<ul> <li>State: Lyophilized purified Ig fraction.</li> <li>Purification: Immunoaffinity chromatography.</li> <li>Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% (w/v) Sodium Azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer.</li> <li>Label: FITC – Fluorescein isothiocyanate (Molecular Weight 390 daltons) Absorption / Emission: 495 nm / 528 nm Molar Ratio: 2.5 moles FITC per mole of Goat IgG.</li> <li>Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).</li> </ul>
Applications:	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Human IgM and Human Serum. No reaction was observed against Human IgG or Human IgA. Specificity was confirmed by ELISA at less than 1% cross reactivity against other human heavy or light chain isotypes.
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.
General References: Conjugation: The and Feltkamp, Immunology 18; 865, 1970.	

**For research and in vitro use only. Not for diagnostic or therapeutic work.** Material Safety Datasheets are available at www.acris-antibodies.com or on request.