

Polyclonal Antibody to Monkey IgA alpha (alpha chain specific) -FITC-

Alternate names:	Monkey Immunoglobulin A
Catalog No.:	R1346F
Quantity:	1 mg
Concentration:	1.0 mg/ml (by UV absorbance at 280 nm)
Host:	Goat
Immunogen:	Monkey IgA α chain.
Format:	State: Lyophilized purified Ig fraction. Purification: Immunoaffinity chromatography. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: FITC – Fluorescein isothiocyanate (Molecular Weight 390 daltons) <i>Absorption / Emission:</i> 495 nm / 528 nm <i>Molar Ratio:</i> 5.0 moles FITC per mole of Goat IgG. Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).
Applications:	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Minimum ELISA Endpoint Dilution: 1:10,000. Actual: 1:40,000. Disposition: PASS. Maximum Cross Reactivity: IgM, IgG < / = 5%. Actual: 3.1% IgM, 1.5% IgG. Disposition: PASS. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Monkey IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Monkey IgA and Monkey Serum. No reaction was observed against Monkey IgM or Monkey IgG.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Storage:

Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use.

Avoid repeated freezing and thawing.

Shelf life: One year from despatch.

General References: Conjugation Reference: The and Feltkamp, Immunology 18; 865, 1970.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com