

Polyclonal Antibody to Rat IgG [H+L] - FITC

Catalog No.: R1371F Quantity: 2 mg

Concentration: 2.0 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Rat IgG whole molecule.

Format: State: Lyophilized purified IgG fraction.

Purification: Immunoaffinity Chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml BSA (lgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as

preservative.

Label: FITC – Fluorescein isothiocyanate (Molecular Weight 390 daltons)

Absorption / Emission: 495 nm / 528 nm

Molar Ratio: 3.1 moles FITC per mole of Rabbit IgG.

Reconstitution: Restore with 1.0 ml of deionized water or equivalent.

Applications: Suitable for Immunomicroscopy and Flow Cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring lot-to-lot consistency.

Recommended Dilution(s): FLISA: 1/10,000-1/50,000. Flow Cytometry: 1/500-1/2,500. Immunofluorescence: 1/1,000-1/5,000.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by Immunoaffinity

Chromatography using Rat IgG coupled to agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities.

Assay by Immunoelectrophoresis resulted in a single precipitin arc against

anti-Fluorescein, anti-Rabbit Serum, Rat IgG and Rat Serum.

Store vial at 2-8°C prior to restoration. Restore with deionized water (or equivalent);

centrifuge product if not completely clear after standing at room temperature. This product

is stable for one month at 2-8°C as an undiluted liquid.

For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Avoid repeated freezing and thawing.

Dilute only prior to immediate use. Shelf life: one year from despatch.

General References: 1. The and Feltkamp, Immunology 18; 865, 1970. (Conjugation)