

## Polyclonal Antibody to Rat IgG F(ab')<sub>2</sub> -Biotin-

<b>Catalog No.:</b>	R1374B
<b>Quantity:</b>	2 mg
<b>Concentration:</b>	2.0 mg/ml (by UV absorbance at 280 nm)
<b>Host:</b>	Rabbit
<b>Immunogen:</b>	Rat IgG F(ab') <sub>2</sub> fragment.
<b>Format:</b>	<b>State:</b> Lyophilized purified Ig fraction. <b>Purification:</b> Immunoaffinity chromatography. <b>Buffer System:</b> 0.01 M Sodium Phosphate, 0.14 M Sodium Chloride, pH 7.4 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.01% (w/v) Thimerisol as preservative. <b>Label:</b> Biotin – Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) <i>Molar Ratio:</i> 10-20 BAC molecules per Rabbit IgG molecule. <b>Reconstitution:</b> Restore with 1.0 ml of deionized water (or equivalent).
<b>Applications:</b>	Immunoblotting, ELISA, Immunohistochemistry, Immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency. <b>Recommended Dilutions:</b> This product has been assayed against 1.0 µg of Rat IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:6,000 to 1:25,000 of the reconstitution concentration is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Rat IgG, Rat IgG F(ab') <sub>2</sub> and Rat Serum. No reaction was observed against Rat IgG F(c).
<b>Storage:</b>	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)

**General References:** 1. Bayer & Wilchek Methods in Enzymology 184; 138-160, 1990.

---

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)