

Polyclonal Antibody to Sheep IgG F(c) -HRP-

Catalog No.: R1380HRP
Quantity: 1.5 mg

Concentration: 1.5 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Sheep IgG F(c) fragment.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.01% (w/v)

Gentamicin Sulfate as preservative. Label: HRP – Horseradish Peroxidase

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunoblotting (Western or Dot blot), ELISA, immunoperoxidase electron

microscopy and Immunohistochemistry as well as other peroxidase-antibody based

enzymatic assays requiring lot-to-lot consistency.

Recommended Dilutions: This product has been assayed against 1.0 ug of Sheep IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000

of the reconstitution concentration is suggested for this product.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Sheep IgG coupled to agarose.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase,

anti-Rabbit Serum, Sheep IgG, Sheep IgG F(c) and Sheep Serum.

No reaction was observed against Sheep IgG F(ab')2.

Storage: Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50%

glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature.

This antibody is stable for one month at 4°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: One year from despatch.

General References: 1. Farr & Nakane, J. Immunol. Methods 47; 129-144. 1981.