

Polyclonal Antibody to Sheep IgG F(c) -AP-

Catalog No.: R1381AP
Quantity: 0.5 mg

Concentration: 1.0 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Sheep IgG F(c) fragment.

Format: State: Liquid (sterile filtered) purified IgG fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.05M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.01% (w/v) sodium azide as preservative. **Label:** AP – Alkaline Phosphatase (Calf Intestine) (Molecular Weight 140,000 daltons)

Applications: Suitable for Immunoblotting (Western or dot blot), ELISA and Immunohistochemistry as

well as other phosphatase-antibody based enzymatic assays requiring lot-to-lot

consistency.

<u>Recommended Dilutions</u>: This product has been assayed against 1.0 ug of Sheep IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:14,000 of the reconstitution

concentration is suggested for this product.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Sheep IgG coupled to agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Sheep IgG, Sheep IgG F(c) and Sheep

Serum.

No reaction was observed against Sheep IgG F(ab')2 or Human Serum Proteins.

Storage: Store the antibody (undiluted) at 2-8°C.

DO NOT FREEZE!

Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic

activity.

Dilute only prior to immediate use. Shelf life: One year from despatch.

General References: 1. Modified from Avarameas and Ternyrock, Immunochemistry 32; 1175 1971.