

Polyclonal Antibody to Sheep IgG F(c) -HRP-

Catalog No.: R1381HRP

Quantity: 0.5 mg

Concentration: 1.0 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Sheep IgG F(c) fragment.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.01% (w/v)

Gentamicin Sulfate as preservative. Label: HRP – Horseradish Peroxidase

Reconstitution: Restore with 0.5 ml of deionized water (or equivalent).

Applications: Suitable for Immunoblotting (Western or dot blot), ELISA, Immunoperoxidase electron

microscopy and Immunohistochemistry as well as other peroxidase-antibody based

enzymatic assays requiring lot-to-lot consistency.

Recommended Dilutions: This product has been assayed against 1.0 µg of Sheep IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to

1:100,000 of the reconstitution concentration is suggested for this product.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Sheep IgG coupled to agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase,

anti-Rabbit Serum, Sheep IgG, Sheep IgG F(c) and Sheep Serum.

No reaction was observed against Sheep IgG F(ab')2 or Human Serum Proteins.

Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50%

glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature.

This antibody is stable for one month at 4°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: One year from despatch.

General References: 1. Farr & Nakane, J. Immunol. Methods 47; 129-144. 1981.