

Polyclonal Antibody to Sheep IgG F(ab')₂ -Biotin-

Catalog No.:	R1383B
Quantity:	0.5 mg
Concentration:	0.5 mg/ml (by UV absorbance at 280 nm)
Host:	Rabbit
Immunogen:	Sheep IgG F(ab') ₂ fragment.
Format:	State: Lyophilized purified IgG fraction. Purification: Immunoaffinity chromatography. Buffer System: in 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: Biotin – N-Hydroxysuccinimide Ester (BAC) <i>Molar Ratio:</i> 10-20 BAC molecules per Rabbit IgG molecule. Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).
Applications:	Suitable for Immunoblotting, ELISA, Immunohistochemistry, Immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency. Recommended Dilutions: This product has been assayed against 1.0 µg of Sheep IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenzothiazoline-6- sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:400,000 is suggested for ELISA. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG F(ab') ₂ coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Sheep IgG and Sheep Serum. No reaction was observed against Human Serum Proteins.
Storage:	Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 4°C as an undiluted liquid. Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: One year from despatch.
General References:	1. Bayer & Wilchek Methods in Enzymology 184; 138-160, 1990.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com