

Polyclonal Antibody to Swine IgG F(ab')2 -HRP-

Alternate names: Pig IgG, Pig Immunoglobulin G

Catalog No.: R1386HRP
Quantity: 2 mg
Host: Rabbit

Immunogen: Swine IgG F(ab')2 fragment.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.01 M Sodium Phosphate, 0.14M Sodium Chloride, pH 7.4 with 0.01% (w/v) Gentamicin sulfate as preservative and 10 mg/ml Bovine Serum Albumin (BSA; IgG

and Protease free) as stabilizer. **Label:** HRP – Horseradish Peroxidase

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunoblotting (Western or Dot blot), ELISA, Immunoperoxidase electron

microscopy and Immunohistochemistry as well as other peroxidase-antibody based

enzymatic assays requiring lot-to-lot consistency.

Recommended Dilutions: This product has been assayed against 1.0 ug of Swine IgG

F(ab')2 in a standard capture ELISA using ABTS

(2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:5,000 to 1:20,000 of the reconstitution

concentration is suggested for this product.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Swine IgG coupled to agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase,

anti-Rabbit Serum, Swine IgG F(ab')2 and Swine Serum.

No reaction was seen against Swine IgG F(c).

Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50%

glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature.

This antibody is stable for one month at 4°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: One year from despatch.

General References: 1. Farr & Nakane, J. Immunol. Methods 47; 129-144. 1981.