

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

R1390TR **Acris Antibodies GmbH** 

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



## Fab Fragment of Goat anti-Rabbit IgG [H&L] -Texas Red-

| Alternate names:   | Rabbit Immunoglobulin G  |
|--|--|
| Catalog No.:   | R1390TR  |
| Quantity:  | 1 mg   |
| Concentration:   | 1.0 mg/ml (by UV absorbance at 280 nm)   |
| Host:  | Goat   |
| Immunogen:   | Rabbit IgG whole molecule.   |
| Format:  | <ul> <li>State: Lyophilized Fab fragments.</li> <li>Purification: Immunoaffinity chromatography.</li> <li>Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative.</li> <li>Label: Texas Red Sulfonyl Chloride (TR; Molecular Weight 625 daltons) Absorption / Emission: 596 nm / 620 nm Molar Ratio: 1.0 moles TR per mole of Goat IgG Fab.</li> <li>Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).</li> </ul> |
| Applications:  | Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other<br>antibody based fluorescent assays requiring lot-to-lot consistency.<br>Other applications not tested. Optimal dilutions are dependent on conditions and should<br>be determined by the user.   |
| Specificity:   | This product was prepared from monospecific antiserum by immunoaffinity<br>chromatography using Rabbit IgG coupled to agarose beads followed by solid phase<br>adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic<br>separation.<br>Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat<br>Serum.  |
|  | No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).  |
| Storage:   | Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below.<br>Centrifuge product if not completely clear after standing at room temperature.<br>This antibody is stable for one month at 4°C as an undiluted liquid.<br>Dilute only prior to immediate use.<br>Avoid cycles of freezing and thawing.<br>Shelf life: One year from despatch.  |
| General References: 1. J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. |  |

General References: 1. J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.