

## F(ab')<sub>2</sub> Fragment of Rabbit anti-Goat IgG (H&L) -FITC -

<b>Alternate names:</b>	Goat Immunoglobulin G
<b>Catalog No.:</b>	R1392F
<b>Quantity:</b>	0.5 mg
<b>Concentration:</b>	1.0 mg/ml (by UV absorbance at 280 nm)
<b>Host:</b>	Rabbit
<b>Immunogen:</b>	Goat IgG whole molecule.
<b>Format:</b>	<b>State:</b> Lyophilized F(ab') <sub>2</sub> fragments. <b>Purification:</b> Immunoaffinity chromatography. <b>Buffer System:</b> 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2 containing 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide , 0.01% (w/v) Thimerosal as preservatives. <b>Label:</b> FITC – Fluorescein isothiocyanate (Molecular Weight 390 daltons) <i>Absorption / Emission:</i> 495 nm / 528 nm <i>Molar Ratio:</i> 3.3 moles FITC per mole of Rabbit IgG F(ab') <sub>2</sub> . <b>Reconstitution:</b> Restore with 0.5 ml of deionized water (or equivalent).
<b>Applications:</b>	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. <b>Recommended Dilution(s):</b> 1:50 to 1:200 for most applications. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Goat IgG and Goat Serum. No reaction was observed against anti-Pepsin, anti-Rabbit IgG F(c) or Human Serum Proteins.
<b>Storage:</b>	Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 4°C as an undiluted liquid. Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: One year from despatch.

**General References:** 1. The and Feltkamp, Immunology 18; 865, 1970.

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)