

F(ab')₂ Fragment of Goat anti-Guinea Pig IgG [H&L] -Texas Red-

| | |
|-------------------------|---|
| Alternate names: | Guinea Pig Immunoglobulin G |
| Catalog No.: | R1393TR |
| Quantity: | 1 mg |
| Concentration: | 1.0 mg/ml (by UV absorbance at 280 nm) |
| Host: | Goat |
| Immunogen: | Guinea Pig IgG whole molecule. |
| Format: | State: Lyophilized F(ab') ₂ fragments. Purification: Immunoaffinity chromatography. Buffer System: 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.6 , containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: Texas Red – -- Sulfonyl Chloride (TR; Molecular Weight 625 daltons) <i>Absorption / Emission:</i> 596 nm / 620 nm <i>Molar Ratio:</i> 3.0 moles Texas Red™ per mole of Goat IgG F(ab') ₂ . Reconstitution: Restore with 1.0 ml of deionized water (or equivalent). |
| Applications: | Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Flow Cytometry: 1:2,000 - 1:10,000. Immunofluorescence microscopy: 1:500 - 1:2,500. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user. |
| Specificity: | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c). |
| Storage: | Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 4°C as an undiluted liquid. Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: One year from despatch. |

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

General References: 1. J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com