

## F(ab')2 Fragment of Goat anti-Human IgG [H&L] -Texas Red-

Alternate names: Human Immunoglobulin G

Catalog No.: R1398TR

Quantity: 1 mg

**Concentration:** 1.0 mg/ml (by UV absorbance at 280 nm)

Host: Goat

Immunogen: Human IgG F(ab')2 fragment.

Format: State: Lyophilized F(ab')2 fragments.

Purification: Immunoaffinity chromatography.

**Buffer System:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v)

Sodium Azide as preservative.

Label: Texas Red - -- Sulfonyl Chloride (TR; Molecular Weight 625 daltons)

Absorption / Emission: 596 nm / 620 nm

*Molar Ratio*: 2.7 moles Texas RedTM per mole of Goat IgG F(ab')2. **Reconstitution**: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other

antibody based fluorescent assays requiring extremely low background levels, absence of

F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

**Specificity:** This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic

separation.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat

Serum, Human IgG, Human IgG F(ab')2 and Human Serum.

No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Human IgG F(c) or Bovine,

Horse, Mouse, Rabbit, Rat and Sheep Serum Proteins.

Storage: Store vial at 4°C prior to restoration. For extended storage add glycerol to 50% and then

aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear

after standing at room temperature.

This antibody is stable for one month at 4°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.



General References: 1. J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982.