

F(ab')₂ Fragment of Rabbit anti-Mouse IgG [H&L] -FITC-

Alternate names:	Mouse Immunoglobulin G
Catalog No.:	R1403F
Quantity:	1 mg
Concentration:	1.0 mg/ml (by UV absorbance at 280 nm)
Host:	Rabbit
Immunogen:	Mouse IgG whole molecule.
Format:	State: Lyophilized F(ab') ₂ fragments. Purification: Immunoaffinity chromatography. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 0.01% (w/v) Sodium Azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer. Label: FITC – Fluorescein isothiocyanate (Molecular Weight 390 daltons) <i>Absorption / Emission:</i> 495 nm / 528 nm <i>Molar Ratio:</i> 2.6 moles FITC per mole of Rabbit IgG F(ab') ₂ . Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).
Applications:	Suitable for Immunomicroscopy and Flow Cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immuno-electrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Rabbit IgG F(c) or Human Serum Proteins.
Storage:	Store vial at 2-8°C prior to restoration. Restore with 1.0 ml of deionized water (or equivalent). For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General References: 1. The and Feltkamp, Immunology 18; 865, 1970.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com