

F(ab)₂ Fragment of Goat anti-Mouse IgA + IgG + IgM (H&L) -Texas Red-

Alternate names:	Mouse Immunoglobulin A
Catalog No.:	R1409TR
Quantity:	1 mg
Concentration:	1.0 mg/ml (by UV absorbance at 280 nm)
Host:	Goat
Immunogen:	Mouse IgG, IgA, and IgM whole molecules.
Format:	State: Lyophilized F(ab') ₂ fragments. Purification: Immunoaffinity chromatography. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% (w/v) Sodium Azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA) as stabilizer. Label: Texas Red – -- Sulfonyl Chloride (Molecular Weight 625 daltons) <i>Absorption / Emission:</i> 596 nm / 620 nm <i>Molar Ratio:</i> 3.0 moles Texas Red TM per of Goat IgG F(ab') ₂ . Reconstitution: Restore with 0.75 ml of deionized water (or equivalent).
Applications:	Suitable for Immunomicroscopy and Flow Cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from polyspecific antiserum by Immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. No reaction was observed against anti-Goat IgG F(c), anti-Pepsin or Human Serum Proteins. This product is suitable for the detection of all Human immunoglobulin classes, isotypes and chain combinations.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Storage:

Store vial at 2-8°C prior to restoration. Restore with 0.75 ml of deionized water (or equivalent). For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use.

Avoid repeated freezing and thawing.

Shelf life: One year from despatch.

General References: 1. J.A. Titus, P.P. Haugland, S.D. Sharrow, D.M. Segal J. Immunol. Methods 50; 193, 1982.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com