

Polyclonal Antibody to Rabbit IgG [H&L] -FITC-

Alternate names: Rabbit Immunoglobulin G

Catalog No.: R1454F Quantity: 1 mg

Concentration: 1.0 mg/ml (by UV absorbance at 280 nm)

Host: Goat

Immunogen: Rabbit IgG, whole molecule.

Format: State: Lyophilized purified IgG fraction.

Purification: Immunoaffinity chromatography using Rabbit IgG coupled to agarose beads

followed by solid phase adsorption(s) to remove any unwanted reactivities

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing

10 mg/ml BSA (IgG and Protease free) and 0.01% (w/v) Sodium Azide. **Label:** FITC – Fluorescein isothiocyanate (Molecular Weight 390 daltons)

Absorption / Emission: 495 nm / 528 nm

Molar Ratio: 3.8 moles FITC per mole of Goat IgG.

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for immunomicroscopy (1:1,000-1:5,000) and flow cytometry or FACS analysis as

well as other antibody based fluorescent assays requiring lot-to-lot consistency.

ELISA: 1/10,000-1/50,000.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: Assay by Immunoelectrophoresis resulted in a single precipitin arc against

anti-Fluorescein, anti-Goat Serum, Rabbit IgG and Rabbit Serum.

No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Horse, Human,

Mouse, Rat or Sheep serum proteins. Actual: all cross reactivities $\langle \ \ | = 0.2\%$.

Storage: Store vial at 2-8°C prior to restoration.

Restore with deionized water (or equivalent); centrifuge product if not completely clear after standing at room temperature. This product is stable for one month at 2-8°C as an undiluted liquid. For extended storage aliquot contents and freeze at -20°C or below.

Avoid repeated freezing and thawing. Dilute only prior to immediate use. Shelf life: one year from despatch.

General References: The and Feltkamp, Immunology 18; 865, 1970. (Conjugation)