

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com R1460TR Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



Polyclonal Antibody to Cyanine Dyes - Texas Red

R1460TR
1 mg
1.0 mg/ml (by UV absorbance at 280 nm)
Sheep
A mixture of Cy3 and Cy5 conjugated KLH.
 State: Lyophilized purified Ig fraction Purification: Immunoaffinity chromatography Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% sodium azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer. Label: Texas Red – TMSulfonyl Chloride (Molecular Weight 625 daltons) Absorption / Emission: 596 nm / 620 nm Molar Ratio: 2.0 moles Texas Red TM per mole of Sheep IgG. Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).
Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
This product was prepared from monospecific antiserum by Immunoaffinity chromatography using a cyanine BSA conjugate coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Sheep Serum and cyanine conjugated IgG. Reactivity was observed against Cy2, Cy3 and Cy5. Reactivity against other cyanine analog is unknown.
Store vial at 2-8°C prior to restoration. For extended storage mix with glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This product is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General References: 1. J. Titus, P. Haugland, S. Sharrow, D. Segal J. Immunol. Methods 50; 193, 1982.