

Polyclonal Antibody to Mouse IgG [H&L] -FITC-

Alternate names: Mouse Immunoglobulin G

Catalog No.: R1611F Quantity: 2 mg

Concentration: 2.0 mg/ml (before lyophil. by UV absorbance at 280 nm)

Host: Goat

Immunogen: Mouse IgG whole molecule.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity Chromatography using Mouse IgG coupled to agarose beads

followed by solid phase adsorption(s) to remove any unwanted reactivities.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 0.01% (w/v) Sodium Azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA, IgG

and Protease free) as stabilizer.

Label: FITC - Fluorescein isothiocyanate; Molecular Weight 390 daltons)

Absorption / Emission: 495 nm / 528 nm Molar Ratio: 3.3 moles FITC per mole of Goat IgG.

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy (1/500-1/2,500) and Flow cytometry or FACS analysis

(1/2,000-1/10,000) as well as other antibody based fluorescent assays requiring lot-to-lot

consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: Assay by immunoelectrophoresis resulted in a single precipitin arc against

anti-Fluorescein, anti-Goat Serum, Mouse IgG and Mouse Serum.

Storage: Store vial at 2-8°C prior to restoration. Restore with deionized water or equivalent.

Centrifuge product if not completely clear after standing at room temperature.

This product is stable for one month at 2-8°C as an undiluted liquid. For extended storage aliquot contents and freeze at -20°C or below. Avoid cycles of freezing and thawing. Dilute only prior to immediate use.

Shelf life: one year from despatch.

General References: The and Feltkamp, Immunology 18; 865, 1970 (Conjugation)